arXiv:1403.6972v1 [math.AC] 27 Mar 2014

Asymptotic prime divisors over complete intersection rings

DIPANKAR GHOSH
TONY J. PUTHENPURAKAL

March 28, 2014

Abstract

Let $(A,\mathfrak{m})$ be a local complete intersection ring. Let M,N be two finitely generated A-modules and I an ideal of A. We prove that

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0}\operatorname{Ass}_A\left(\operatorname{Ext}^i_A(M,N/I^nN)\right)$$

is a finite set. Moreover, we prove that there exists $i_0,n_0\geq 0$ such that for all $i\geq i_0$ and $n\geq n_0$, we have

$$\operatorname{Ass}_A\left(\operatorname{Ext}^{2i}_A(M,N/I^nN)\right)=\operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0}_A(M,N/I^{n_0}N)\right),$$
$$\operatorname{Ass}_A\left(\operatorname{Ext}^{2i+1}_A(M,N/I^nN)\right)=\operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0+1}_A(M,N/I^{n_0}N)\right).$$

We also prove the analogous result for complete intersection rings which arise in algebraic geometry. Further, we prove that the complexity $\operatorname{cx}_A(M,N/I^nN)$ is constant for all sufficiently large n.

1 Introduction

Let A be a commutative Noetherian ring, I an ideal of A, and M a finitely generated A-module. Brodmann [2] proved that the set of associated prime ideals $\operatorname{Ass}_A(M/I^nM)$ is independent of n for all sufficiently large n. Thereafter, L. Melkersson and P. Schenzel generalized Brodmann's result in [6, Theorem 1] by proving that

$$\operatorname{Ass}_A\left(\operatorname{Tor}^A_i(M,A/I^n)\right)$$

is independent of n for all large n and for a fixed $i\geq 0$.

Later, D. Katz and E. West proved the above result in a more general way [5, 3.5]; if N is a finitely generated A-module, then for a fixed $i\geq 0$, the sets

$$\operatorname{Ass}_A\left(\operatorname{Tor}^A_i(M,N/I^nN)\right)\quad\text{and}\quad\operatorname{Ass}_A\left(\operatorname{Ext}^i_A(M,N/I^nN)\right)$$

are stable for all large n. So, in particular, for a fixed $i \geq 0$,

$$\bigcup_{n\geq 0} \operatorname{Ass}_A \left(\operatorname{Ext}^i_A(M, N/I^n N)\right)$$

is a finite set. In this context, the second author [7, page 368] raised a question about what happens if we vary i (≥ 0) also? More precisely,

(†) is the set $\bigcup_{i\geq 0}\bigcup_{n\geq 0} \operatorname{Ass}_A \left(\operatorname{Ext}^i_A(M, N/I^n N)\right)$ finite?

The motivation for the question (†) came from the following two questions. They were raised by W. Vasconcelos [8, 3.5] and Melkersson and Schenzel [6, page 936] respectively.

(1) Is the set $\bigcup_{i\geq 0} \operatorname{Ass}_A \left(\operatorname{Ext}^i_A(M, A)\right)$ finite?

(2) Is the set $\bigcup_{i\geq 0}\bigcup_{n\geq 1} \operatorname{Ass}_A \left(\operatorname{Tor}^A_i(M, A/I^n)\right)$ finite?

Recently, the second author [7, Theorem 5.1] proved that if A is a local complete intersection ring and $\mathcal{N} = \oplus_{n\geq 0} N_n$ is a finitely generated graded $\mathscr{R}(I)$-module, then

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0} \operatorname{Ass}_A \left(\operatorname{Ext}^i_A(M, N_n)\right)$$

is a finite set. Moreover, he proved that there exists $i_0, n_0 \geq 0$ such that

$$\begin{aligned}\operatorname{Ass}_A \left(\operatorname{Ext}^{2i}_A(M, N_n)\right) &= \operatorname{Ass}_A \left(\operatorname{Ext}^{2i_0}_A(M, N_{n_0})\right),\\ \operatorname{Ass}_A \left(\operatorname{Ext}^{2i+1}_A(M, N_n)\right) &= \operatorname{Ass}_A \left(\operatorname{Ext}^{2i_0+1}_A(M, N_{n_0})\right)\end{aligned}$$

for all $i \geq i_0$ and $n \geq n_0$. In particular, if N is a finitely generated A-module, then $\mathcal{N}$ can be taken as $\bigoplus_{n\geq 0}(I^n N)$ or $\bigoplus_{n\geq 0}(I^n N/I^{n+1}N)$. In the present study, we prove that the question (†) has an affirmative answer for a local complete intersection ring. We also analyze the stability of the sets of associated prime ideals which occurs periodically after certain stage.

Let $(A, \mathfrak{m})$ be a local complete intersection ring. Let M, N be two finitely generated A-modules and I an ideal of A. Define

$$\operatorname{cx}_A(M, N) = \inf\left\{b \in \mathbb{N} \;\middle|\; \limsup_{n\to\infty} \frac{\mu(\operatorname{Ext}^n_A(M, N))}{n^{b-1}} < \infty\right\},$$

where $\mu(D)$ denote the number of minimal generators of a finitely generated A-module D. In [7, Theorem 7.1], the second author proved that $\operatorname{cx}_A(M, I^j N)$ is constant for all $j \gg 0$. In Section 6, we prove that

(††) $\operatorname{cx}_A(M, N/I^j N)$ is constant for all $j \gg 0$.

The rest of the paper are organized as follows. In Section 2, we give some graded module structures which we use in order to prove our main results. The finiteness result on asymptotic primes are proved in Section 3; while the stability result is shown in Section 4. Then, in Section 5, we prove the analogous results on asymptotic primes for complete intersection rings which arise in algebraic geometry. Finally, in Section 6, we prove (††).

2 Module structure

In this section, we give the graded module structures which we are going to use in order to prove our main results.

Let Q be a commutative Noetherian ring and $\mathbf{f} = f_1, \ldots, f_c$ a Q-regular sequence. Set $A = Q/(\mathbf{f})$. Let M, D be two A-modules and let M be finitely generated.

2.1. Let $\mathbb{F}: \cdots \to F_n \to \cdots \to F_1 \to F_0 \to 0$ be a projective resolution of M by finitely generated free A-modules. Let $t_j : \mathbb{F}(+2) \to \mathbb{F}$, $1 \leq j \leq c$ be the Eisenbud operators defined by $\mathbf{f} = f_1, \ldots, f_c$ (see [3, Section 1]). By [3, 1.4], the maps t_j are determined uniquely up to homotopy. In particular, they induce well defined maps

$$t_j : \operatorname{Ext}^i_A(M, D) \to \operatorname{Ext}^{i+2}_A(M, D) \quad \text{for all } i \text{ and } 1 \leq j \leq c.$$

By [3, 1.5], the maps t_j $(j = 1, \ldots, c)$ commute up to homotopy. Thus

$$\operatorname{Ext}^\star_A(M, D) := \bigoplus_{i \geq 0} \operatorname{Ext}^i_A(M, D)$$

turns into a graded $T := A[t_1, \ldots, t_c]$-module where T is the graded polynomial ring over A in the cohomology operators t_j defined by $\mathbf{f}$ with $\deg(t_j) = 2$ for all $1 \leq j \leq c$. Furthermore, these structures depend only on $\mathbf{f}$, are natural in both module arguments and commute with the connecting maps induced by short exact sequences.

2.2. If $\operatorname{projdim}_Q(M)$ is finite and D is a finitely generated A-module, then Gulliksen [4, 3.1] proved that $\operatorname{Ext}^\star_A(M, D)$ is a finitely generated graded T-module.

2.3. Let I be an ideal of A. Let $\mathscr{R}(I) = \oplus_{n \geq 0} I^n X^n$ be the Rees ring associated to I. We consider $\mathscr{R}(I)$ as a subring of the polynomial ring $A[X]$. Let $\mathcal{N} = \oplus_{n \geq 0} N_n$ be a graded $\mathscr{R}(I)$-module. Let $u \in \mathscr{R}(I)$ be a homogeneous element of degree s. Consider the A-linear maps given by multiplication with u:

$$N_n \xrightarrow{u} N_{n+s} \quad \text{for all } n.$$

By applying $\operatorname{Hom}_A(\mathbb{F}, -)$ on the above maps and using the naturality of the operators t_j, we have the following commutative diagram of complexes:

$$\begin{array}{ccc} \operatorname{Hom}_A(\mathbb{F}, N_n) & \xrightarrow{t_j} & \operatorname{Hom}_A(\mathbb{F}(+2), N_n) \\ \downarrow u & & \downarrow u \\ \operatorname{Hom}_A(\mathbb{F}, N_{n+s}) & \xrightarrow{t_j} & \operatorname{Hom}_A(\mathbb{F}(+2), N_{n+s}). \end{array}$$

Now, taking cohomology, we have the following commutative diagram of A-modules:

$$\begin{array}{ccc} \mathrm{Ext}^i_A(M, N_n) & \xrightarrow{t_j} & \mathrm{Ext}^{i+2}_A(M, N_n) \\ \downarrow u & & \downarrow u \\ \mathrm{Ext}^i_A(M, N_{n+s}) & \xrightarrow{t_j} & \mathrm{Ext}^{i+2}_A(M, N_{n+s}) \end{array}$$

for all i, n and $1 \leq j \leq c$. Thus

$$\mathscr{E}(\mathcal{N}) := \bigoplus_{i\geq 0} \bigoplus_{n\geq 0} \mathrm{Ext}^i_A(M, N_n)$$

turns into a bigraded $\mathscr{S} := \mathscr{R}(I)[t_1, \ldots, t_c]$-module where we set $\deg(t_j) = (0, 2)$ for all $1 \leq j \leq c$ and $\deg(uX^s) = (s, 0)$ for all $u \in I^s$, $s \geq 0$.

2.4. Suppose N is a finitely generated A-module. Set $\mathcal{L} := \oplus_{n\geq 0}(N/I^{n+1}N)$. Note that $\mathscr{R}(I, N) = \oplus_{n\geq 0} I^n N$ and $N[X] = N \otimes_A A[X]$ are graded modules over $\mathscr{R}(I)$ and $A[X]$ respectively. Since $\mathscr{R}(I)$ is a graded subring of $A[X]$, we set that $N[X]$ is a graded $\mathscr{R}(I)$-module. Therefore $\mathcal{L}$ is a graded $\mathscr{R}(I)$-module where the graded structure is induced by the sequence

$$0 \to \mathscr{R}(I, N) \to N[X] \to \mathcal{L}(-1) \to 0.$$

Therefore, from Section 2.3,

$$\mathscr{E}(\mathcal{L}) = \bigoplus_{i\geq 0} \bigoplus_{n\geq 0} \mathrm{Ext}^i_A(M, N/I^{n+1}N)$$

is a bigraded module over $\mathscr{S} = \mathscr{R}(I)[t_1, \ldots, t_c]$.

Let us recall the following result from [7, 1.1].

Theorem 2.5. *Let Q be a Noetherian ring with finite Krull dimension and let $\mathbf{f} = f_1, \ldots, f_c$ be a regular sequence in Q. Set $A = Q/(\mathbf{f})$. Let M be a finitely generated A-module with $\mathrm{projdim}_Q(M)$ finite. Let I be an ideal of A and let $\mathcal{N} = \oplus_{n\geq 0} N_n$ be a finitely generated $\mathscr{R}(I)$-module. Then*

$$\mathscr{E}(\mathcal{N}) := \bigoplus_{i\geq 0} \bigoplus_{n\geq 0} \mathrm{Ext}^i_A(M, N_n)$$

is a finitely generated bigraded $\mathscr{S} = \mathscr{R}(I)[t_1, \ldots, t_c]$-module.

We first prove the main result of this article for a ring A which is of the form $Q/(\mathbf{f})$, where Q is a regular local ring and $\mathbf{f} = f_1, \ldots, f_c$ is a Q-regular sequence. Then we deduce the main result for a local complete intersection ring with the help of the following lemma:

Lemma 2.6. ([7, 5.6.(b)]) *Let $(A, \mathfrak{m})$ be a Noetherian local ring and $\widehat{A}$ the completion of A with respect to $\mathfrak{m}$. Let D be an A-module. Then*

$$\mathrm{Ass}_A(D) = \{\mathfrak{q} \cap A : \mathfrak{q} \in \mathrm{Ass}_{\widehat{A}}(D \otimes_A \widehat{A})\}.$$

3 Asymptotic associated primes: Finiteness

In this section, we prove the finiteness (Corollary 3.2) of the set of asymptotic associated primes of a family of Ext modules over a local complete intersection ring.

Theorem 3.1. *Let Q be a Noetherian ring with finite Krull dimension, and $\mathbf{f} = f_1, \ldots, f_c$ a Q-regular sequence. Set $A = Q/(\mathbf{f})$. Let M, N be two finitely generated A-modules with $\operatorname{projdim}_Q(M)$ finite, and I an ideal of A. Then*

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0} \operatorname{Ass}_A\left(\operatorname{Ext}^i_A(M, N/I^nN)\right) \quad \textit{is a finite set.}$$

Proof. For a fixed $n \geq 0$, consider the short exact sequence of A-modules:

$$0 \longrightarrow I^nN/I^{n+1}N \longrightarrow N/I^{n+1}N \longrightarrow N/I^nN \longrightarrow 0.$$

Taking direct sum over $n \geq 0$, and setting $\mathcal{L} := \oplus_{n\geq 0}(N/I^{n+1}N)$, we have a short exact sequence of graded $\mathscr{R}(I)$-modules:

$$0 \longrightarrow \operatorname{gr}_I(N) \longrightarrow \mathcal{L} \longrightarrow \mathcal{L}(-1) \longrightarrow 0,$$

which induces an exact sequence of $\mathscr{R}(I)$-modules for each $i \geq 0$:

$$\operatorname{Ext}^i_A(M, \operatorname{gr}_I(N)) \longrightarrow \operatorname{Ext}^i_A(M, \mathcal{L}) \longrightarrow \operatorname{Ext}^i_A(M, \mathcal{L}(-1)).$$

Taking direct sum over $i \geq 0$ and using the naturality of the cohomology operators t_j, we have an exact sequence of $\mathscr{S} = \mathscr{R}(I)[t_1, \ldots, t_c]$-modules:

$$\bigoplus_{i,n\geq 0} \operatorname{Ext}^i_A\left(M, \frac{I^nN}{I^{n+1}N}\right) \xrightarrow{\Phi} \bigoplus_{i,n\geq 0} V_{i,n} \xrightarrow{\Psi} \bigoplus_{i,n\geq 0} V_{i,n-1},$$

where $V_{i,n} := \operatorname{Ext}^i_A(M, N/I^{n+1}N)$ for each $i \geq 0, n \geq -1$. Let

$$U = \bigoplus_{i,n\geq 0} U_{i,n} := \operatorname{Image}(\Phi).$$

Then for each $i, n \geq 0$, considering the exact sequence of A-modules:

$$0 \to U_{i,n} \to V_{i,n} \to V_{i,n-1},$$

we have

$$\begin{aligned}
\operatorname{Ass}_A(V_{i,n}) &\subseteq \operatorname{Ass}_A(U_{i,n}) \cup \operatorname{Ass}_A(V_{i,n-1}) \\
&\subseteq \operatorname{Ass}_A(U_{i,n}) \cup \operatorname{Ass}_A(U_{i,n-1}) \cup \operatorname{Ass}_A(V_{i,n-2}) \\
&\vdots \\
&\subseteq \bigcup_{0\leq j\leq n} \operatorname{Ass}_A(U_{i,j}). \quad [\text{Since } \operatorname{Ass}_A(V_{i,-1}) = \phi \text{ for each } i \geq 0].
\end{aligned}$$

Taking union over $i, n \geq 0$, we have

$$\bigcup_{i,n\geq 0} \operatorname{Ass}_A(V_{i,n}) \subseteq \bigcup_{i,n\geq 0} \operatorname{Ass}_A(U_{i,n}). \tag{1}$$

Since $\operatorname{gr}_I(N)$ is a finitely generated graded $\mathscr{R}(I)$-module, by Theorem 2.5,

$$\bigoplus_{i,n\geq 0} \operatorname{Ext}^i_A\left(M, \frac{I^n N}{I^{n+1}N}\right)$$

is a finitely generated bigraded $\mathscr{S}$-module, and hence U is a finitely generated bigraded $\mathscr{S}$-module. Therefore by [9, Lemma 3.2],

$$\bigcup_{i,n\geq 0} \operatorname{Ass}_A(U_{i,n}) \quad \text{is a finite set.} \tag{2}$$

The result follows from (1) and (2). □

An immediate corollary is the following:

Corollary 3.2. *Let $(A, \mathfrak{m})$ be a local complete intersection ring. Let M, N be two finitely generated A-modules and I an ideal of A. Then*

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0} \operatorname{Ass}_A\left(\operatorname{Ext}^i_A(M, N/I^n N)\right) \quad \textit{is a finite set.}$$

Proof. Since A is a local complete intersection ring, $\widehat{A} = Q/(\mathbf{f})$, where Q is a regular local ring and $\mathbf{f} = f_1, \ldots, f_c$ is a Q-regular sequence.

Applying Theorem 3.1 for the ring $\widehat{A}$, we have

$$\bigcup_{i,n\geq 0} \operatorname{Ass}_{\widehat{A}}\left(\operatorname{Ext}^i_A(M, N/I^n N) \otimes_A \widehat{A}\right) = \bigcup_{i,n\geq 0} \operatorname{Ass}_{\widehat{A}}\left(\operatorname{Ext}^i_{\widehat{A}}\left(\widehat{M}, \widehat{N}/(I\widehat{A})^n \widehat{N}\right)\right)$$

is a finite set, and hence the result follows from Lemma 2.6. □

4 Asymptotic associated primes: Stability

In this section, we prove the stability result (Corollary 4.3). We first prove the following theorem:

Theorem 4.1. *Let Q be a Noetherian ring with finite Krull dimension, and $\mathbf{f} = f_1, \ldots, f_c$ a Q-regular sequence. Set $A = Q/(\mathbf{f})$. Let M, N be two finitely generated A-modules with $\operatorname{projdim}_Q(M)$ finite, and I an ideal of A. Then there exists $i_0, n_0 \geq 0$ such that for all $i \geq i_0$ and $n \geq n_0$, we have*

$$\operatorname{Ass}_A\left(\operatorname{Ext}^{2i}_A(M, N/I^n N)\right) = \operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0}_A(M, N/I^{n_0} N)\right),$$
$$\operatorname{Ass}_A\left(\operatorname{Ext}^{2i+1}_A(M, N/I^n N)\right) = \operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0+1}_A(M, N/I^{n_0} N)\right).$$

To prove this result, we assume the following lemma which we prove at the end of this section.

Lemma 4.2. *Let $(Q, \mathfrak{n})$ be a Noetherian local ring with residue field k, and let $\mathbf{f} = f_1, \ldots, f_c$ be a Q-regular sequence. Set $A = Q/(\mathbf{f})$. Let M, N be two finitely generated A-modules with $\operatorname{projdim}_Q(M)$ finite, and I an ideal of A. Then*

$$\lambda_A\left(\operatorname{Hom}_A\left(k, \operatorname{Ext}_A^{2i}(M, N/I^nN)\right)\right) \quad \text{and} \quad \lambda_A\left(\operatorname{Hom}_A\left(k, \operatorname{Ext}_A^{2i+1}(M, N/I^nN)\right)\right)$$

are given by polynomials in i, n with rational coefficients for all large (i, n).

Proof of Theorem 4.1. Applying Theorem 3.1, we may assume that

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0} \operatorname{Ass}_A\left(\operatorname{Ext}_A^i(M, N/I^nN)\right) = \{\mathfrak{p}_1, \mathfrak{p}_2, \ldots, \mathfrak{p}_l\}.$$

Let $V_{i,n} := \operatorname{Ext}_A^i(M, N/I^nN)$ for each $i, n \geq 0$, and $V := \oplus_{i,n\geq 0} V_{i,n}$.

We first prove that there exists some $i', n' \geq 0$ such that

$$\operatorname{Ass}_A(V_{2i,n}) = \operatorname{Ass}_A(V_{2i',n'}) \quad \text{for all } i \geq i', n \geq n'. \tag{3}$$

To prove the result (3), it is enough to prove that for each $\mathfrak{p}_j$, where $1 \leq j \leq l$, there exists some $i_{j_0}, n_{j_0} \geq 0$ such that exactly one of the following alternatives must hold:

$$\begin{aligned} \text{either} \quad & \mathfrak{p}_j \in \operatorname{Ass}_A(V_{2i,n}) \quad \text{for all } i \geq i_{j_0}, n \geq n_{j_0}; \\ \text{or} \quad & \mathfrak{p}_j \notin \operatorname{Ass}_A(V_{2i,n}) \quad \text{for all } i \geq i_{j_0}, n \geq n_{j_0}. \end{aligned}$$

Localizing at $\mathfrak{p}_j$, and replacing $A_{\mathfrak{p}_j}$ by A and $\mathfrak{p}_j A_{\mathfrak{p}_j}$ by $\mathfrak{m}$, it is now enough to prove that there exists $i', n' \geq 0$ such that

$$\text{either} \quad \mathfrak{m} \in \operatorname{Ass}_A(V_{2i,n}) \quad \text{for all } i \geq i', n \geq n'; \tag{4}$$

$$\text{or} \quad \mathfrak{m} \notin \operatorname{Ass}_A(V_{2i,n}) \quad \text{for all } i \geq i', n \geq n'. \tag{5}$$

But from Lemma 4.2, $\lambda_A(\operatorname{Hom}_A(k, V_{2i,n}))$ is given by a polynomial in i, n with rational coefficients for all large (i, n), and hence there exists $i', n' \geq 0$ such that

$$\begin{aligned} \text{either} \quad & \operatorname{Hom}_A(k, V_{2i,n}) \neq 0 \quad \text{for all } i \geq i', n \geq n'; \\ \text{or} \quad & \operatorname{Hom}_A(k, V_{2i,n}) = 0 \quad \text{for all } i \geq i', n \geq n', \end{aligned}$$

which is equivalent to that either (4) is true, or (5) is true.

Applying similar procedure as in the even case, there exists $i'', n'' \geq 0$ such that

$$\operatorname{Ass}_A(V_{2i+1,n}) = \operatorname{Ass}_A(V_{2i''+1,n''}) \quad \text{for all } i \geq i'', n \geq n''.$$

Then $(i_0, n_0) := \max\{(i', n'), (i'', n'')\}$ satisfies the required result of the theorem. □

An immediate corollary of the Theorem 4.1 is the following:

Corollary 4.3. *Let $(A, \mathfrak{m})$ be a local complete intersection ring. Let M, N be two finitely generated A-modules and I an ideal of A. Then there exists $i_0, n_0 \geq 0$ such that for all $i \geq i_0$ and $n \geq n_0$, we have*

$$\begin{aligned}\mathrm{Ass}_A\left(\mathrm{Ext}_A^{2i}(M, N/I^nN)\right) &= \mathrm{Ass}_A\left(\mathrm{Ext}_A^{2i_0}(M, N/I^{n_0}N)\right),\\ \mathrm{Ass}_A\left(\mathrm{Ext}_A^{2i+1}(M, N/I^nN)\right) &= \mathrm{Ass}_A\left(\mathrm{Ext}_A^{2i_0+1}(M, N/I^{n_0}N)\right).\end{aligned}$$

Proof. Assume $\widehat{A} = Q/(\mathbf{f})$, where Q is a regular local ring and $\mathbf{f} = f_1, \ldots, f_c$ is a Q-regular sequence. Then by applying Theorem 4.1 for the ring $\widehat{A}$, there exists $i_0, n_0 \geq 0$ such that for all $i \geq i_0$ and $n \geq n_0$, we have

$$\begin{aligned}\mathrm{Ass}_{\widehat{A}}\left(\mathrm{Ext}_A^{2i}(M, N/I^nN) \otimes_A \widehat{A}\right) &= \mathrm{Ass}_{\widehat{A}}\left(\mathrm{Ext}_A^{2i_0}(M, N/I^{n_0}N) \otimes_A \widehat{A}\right),\\ \mathrm{Ass}_{\widehat{A}}\left(\mathrm{Ext}_A^{2i+1}(M, N/I^nN) \otimes_A \widehat{A}\right) &= \mathrm{Ass}_{\widehat{A}}\left(\mathrm{Ext}_A^{2i_0+1}(M, N/I^{n_0}N) \otimes_A \widehat{A}\right).\end{aligned}$$

The result now follows from Lemma 2.6. □

We now give

Proof of Lemma 4.2. For each fixed $n \geq 0$, consider the short exact sequence of A-modules:

$$0 \longrightarrow I^nN \longrightarrow N \longrightarrow N/I^nN \longrightarrow 0,$$

which induces an exact sequence of A-modules for each i, n:

$$\mathrm{Ext}_A^i(M, I^nN) \to \mathrm{Ext}_A^i(M, N) \to \mathrm{Ext}_A^i(M, N/I^nN) \to \mathrm{Ext}_A^{i+1}(M, I^nN).$$

Taking direct sum over i, n and using the naturality of the cohomology operators t_j, we have an exact sequence of $\mathscr{S} = \mathscr{R}(I)[t_1, \ldots, t_c]$-modules:

$$U \longrightarrow T \longrightarrow V \longrightarrow U(0,1),$$

where

$$\begin{aligned}U &= \bigoplus_{i,n\geq 0} U_{i,n} := \bigoplus_{i,n\geq 0} \mathrm{Ext}_A^i(M, I^nN),\\ T &= \bigoplus_{i,n\geq 0} T_{i,n} := \bigoplus_{i,n\geq 0} \mathrm{Ext}_A^i(M, N),\\ V &= \bigoplus_{i,n\geq 0} V_{i,n} := \bigoplus_{i,n\geq 0} \mathrm{Ext}_A^i(M, N/I^nN), \quad \text{and}\end{aligned}$$

$U(0,1)$ is same as U but the grading is twisted by $(0,1)$. Setting $X := \mathrm{Image}(U \to T)$, $Y := \mathrm{Image}(T \to V)$ and $Z := \mathrm{Image}(V \to U(0,1))$, we have the following commutative diagram of exact sequences of $\mathscr{S}$-modules:



which gives the following two short exact sequences of $\mathscr{S}$-modules:

$$0 \to X \to T \to Y \to 0 \quad \text{and} \quad 0 \to Y \to V \to Z \to 0.$$

Now applying $\mathrm{Hom}_A(k, -)$ to the above short exact sequences, we get the following exact sequences of $\mathscr{S}$-modules:

$$0 \to \mathrm{Hom}_A(k, X) \to \mathrm{Hom}_A(k, T) \to \mathrm{Hom}_A(k, Y) \to C \to 0, \tag{6}$$

$$0 \to \mathrm{Hom}_A(k, Y) \to \mathrm{Hom}_A(k, V) \to D \to 0, \tag{7}$$

where $C := \mathrm{Image}(\mathrm{Hom}_A(k, Y) \to \mathrm{Ext}^1_A(k, X))$,
$D := \mathrm{Image}(\mathrm{Hom}_A(k, V) \to \mathrm{Hom}_A(k, Z))$.

By Theorem 2.5, $U = \oplus_{i,n\geq 0}\mathrm{Ext}^i_A(M, I^n N)$ is a finitely generated bigraded $\mathscr{S}$-module, hence $X = \mathrm{Image}(U \to T)$ is so, and hence $\mathrm{Hom}_A(k, X)$, $\mathrm{Ext}^1_A(k, X)$ are also finitely generated $\mathscr{S}$-modules. Being an $\mathscr{S}$-submodule of $\mathrm{Ext}^1_A(k, X)$, C is also a finitely generated bigraded $\mathscr{S}$-module. Since $\mathrm{Hom}_A(k, X)$ and $\mathrm{Ext}^1_A(k, X)$ are annihilated by the maximal ideal of A, $\mathrm{Hom}_A(k, X_{i,n})$ and $C_{i,n}$ both are finitely generated k-modules, and hence they have finite length as A-modules for each $i, n \geq 0$. Therefore applying Hilbert-Serre theorem on the $\mathscr{S}$-modules $\mathrm{Hom}_A(k, X)$ and C, we get

$$\sum_{i,n\geq 0} \lambda_A(\mathrm{Hom}_A(k, X_{i,n}))z^i w^n = \frac{P_X(z, w)}{(1 - z^2)^c(1 - w)^r}, \tag{8}$$

$$\sum_{i,n\geq 0} \lambda_A(C_{i,n})z^i w^n = \frac{P_C(z, w)}{(1 - z^2)^c(1 - w)^r} \tag{9}$$

for some polynomials $P_X(z, w)$ and $P_C(z, w)$ over $\mathbb{Z}$.

For a fixed $n \geq 0$, $\oplus_{i\geq 0}T_{i,n} = \oplus_{i\geq 0}\mathrm{Ext}^i_A(M, N)$ is a finitely generated graded $A[t_1, \ldots, t_c]$-module, and hence $\mathrm{Hom}_A(k, \oplus_{i\geq 0}T_{i,n})$ is also so. By similar argument as before $\lambda_A(\mathrm{Hom}_A(k, T_{i,n})) < \infty$ for each $i \geq 0$. Therefore again by Hilbert-Serre theorem, we have for each $n \geq 0$:

$$\sum_{i\geq 0} \lambda_A(\mathrm{Hom}_A(k, T_{i,n}))z^i = \frac{P_S(z)}{(1 - z^2)^c}$$

for some polynomial $P_S(z) \in \mathbb{Z}[z]$. Multiplying both sides of the above equation by w^n, and then taking sum over $n \geq 0$, we have

$$\sum_{i,n\geq 0} \lambda_A(\mathrm{Hom}_A(k, T_{i,n}))z^i w^n = \frac{P_S(z)}{(1 - z^2)^c(1 - w)} = \frac{P_T(z, w)}{(1 - z^2)^c(1 - w)^r}, \tag{10}$$

where $P_T(z, w) = P_S(z)(1 - w)^{r-1} \in \mathbb{Z}[z, w]$.

Now considering (6), we have an exact sequence of A-modules:

$$0 \to \mathrm{Hom}_A(k, X_{i,n}) \to \mathrm{Hom}_A(k, T_{i,n}) \to \mathrm{Hom}_A(k, Y_{i,n}) \to C_{i,n} \to 0$$

for each $i, n \geq 0$. Then additive property of length function gives

$$\lambda_A(\mathrm{Hom}_A(k, X_{i,n})) - \lambda_A(\mathrm{Hom}_A(k, T_{i,n})) + \lambda_A(\mathrm{Hom}_A(k, Y_{i,n})) - \lambda_A(C_{i,n}) = 0$$

for each $i, n \geq 0$. Multiplying the above equation by $z^i w^n$, then taking sum over $i, n \geq 0$, and using (8), (9) and (10), we get

$$\sum_{i,n\geq 0} \lambda_A(\mathrm{Hom}_A(k, Y_{i,n}))z^i w^n = \frac{P_Y(z,w)}{(1-z^2)^c(1-w)^r}, \tag{11}$$

where $P_Y(z,w) = -P_X(z,w) + P_T(z,w) + P_C(z,w) \in \mathbb{Z}[z,w]$.

Recall that by Theorem 2.5, U is a finitely generated bigraded $\mathscr{S}$-module. Now observe that Z is a submodule of $U(0,1)$. Therefore Z is a finitely generated bigraded $\mathscr{S}$-module, hence $\mathrm{Hom}_A(k, Z)$ is so, and hence

$$D = \mathrm{Image}(\mathrm{Hom}_A(k, V) \to \mathrm{Hom}_A(k, Z))$$

is also so. Observe that $\lambda_A(D_{i,n}) < \infty$ for each $i, n \geq 0$. Therefore once again by applying Hilbert-Serre theorem, we have

$$\sum_{i,n\geq 0} \lambda_A(D_{i,n})z^i w^n = \frac{P_D(z,w)}{(1-z^2)^c(1-w)^r} \tag{12}$$

for some polynomial $P_D(z,w)$ over $\mathbb{Z}$.

Next considering (7), we have an exact sequence of A-modules:

$$0 \to \mathrm{Hom}_A(k, Y_{i,n}) \to \mathrm{Hom}_A(k, V_{i,n}) \to D_{i,n} \to 0$$

for each $i, n \geq 0$, which gives

$$\lambda_A(\mathrm{Hom}_A(k, Y_{i,n})) - \lambda_A(\mathrm{Hom}_A(k, V_{i,n})) + \lambda_A(D_{i,n}) = 0.$$

Multiplying the above equation by $z^i w^n$, then taking sum over $i, n \geq 0$, and using (11) and (12), we get

$$\sum_{i,n\geq 0} \lambda_A(\mathrm{Hom}_A(k, V_{i,n}))z^i w^n = \frac{P_V(z,w)}{(1-z^2)^c(1-w)^r},$$

where $P_V(z,w) = P_Y(z,w) + P_D(z,w) \in \mathbb{Z}[z,w]$. Therefore it follows that

$$\lambda_A(\mathrm{Hom}_A(k, V_{2i,n})) \quad \text{and} \quad \lambda_A(\mathrm{Hom}_A(k, V_{2i+1,n}))$$

are given by polynomials in i, n with rational coefficients for all sufficiently large (i, n), which completes the proof of the lemma. □

5 Asymptotic associated primes: The geometric case

Let V be an affine or projective variety over an algebraically closed field K. Let A be the coordinate ring of V. Then V is said to be a locally complete intersection variety if all its local rings are complete intersection. Thus

- in the affine case, $A_{\mathfrak{p}}$ is a local complete intersection ring for every $\mathfrak{p} \in \operatorname{Spec}(A)$.
- in the projective case, $A_{(\mathfrak{p})}$ is a local complete intersection ring for every $\mathfrak{p} \in \operatorname{Proj}(A)$. Recall that $A_{(\mathfrak{p})}$ is the degree zero part of the graded ring $S^{-1}A$, where S is the collection of all homogeneous elements in $A \setminus \mathfrak{p}$.

In this section, we prove the results analogous to Theorems 3.1 and 4.1 for the coordinate rings of locally complete intersection varieties.

In the affine case, we prove the following general result:

Theorem 5.1. *Let Q be a regular ring of finite Krull dimension and $\mathfrak{a}$ an ideal of Q. Set $A = Q/\mathfrak{a}$. Let A be a complete intersection ring. Let M, N be two finitely generated A-modules and I an ideal of A. Then*

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0} \operatorname{Ass}_A\left(\operatorname{Ext}^i_A(M, N/I^nN)\right) \quad \textit{is a finite set.}$$

Moreover, there exists $i_0, n_0 \geq 0$ such that for all $i \geq i_0, n \geq n_0$, we have

$$\operatorname{Ass}_A\left(\operatorname{Ext}^{2i}_A(M, N/I^nN)\right) = \operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0}_A(M, N/I^{n_0}N)\right),$$
$$\operatorname{Ass}_A\left(\operatorname{Ext}^{2i+1}_A(M, N/I^nN)\right) = \operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0+1}_A(M, N/I^{n_0}N)\right).$$

Proof. For $x \in A$, set $D(x) = \{\mathfrak{p} \in \operatorname{Spec}(A) : x \notin \mathfrak{p}\}$. As in [7, *Proof of Theorem 6.1*], we have

$$\operatorname{Spec}(A) = D(g_1) \cup \cdots \cup D(g_m) \quad \text{for some } g_1, \ldots, g_m \in A,$$

such that the localization A_{g_j} at $\{{g_j}^l : l \geq 0\}$ has the form $Q_j/\mathfrak{a}_j$ for some regular ring Q_j of finite Krull dimension and some ideal $\mathfrak{a}_j$ of Q_j generated by a Q_j-regular sequence. Then clearly for any A-module E, we have

$$\operatorname{Ass}_A(E) = \bigcup\{\mathfrak{q} \cap A : \mathfrak{q} \in \operatorname{Ass}_{A_{g_j}}(E_{g_j}) \text{ for some } j = 1, \ldots, m\}. \tag{13}$$

Since localization A_{g_j} is flat over A, we have

$$\left(\operatorname{Ext}^i_A(M, N/I^nN)\right)_{g_j} = \operatorname{Ext}^i_{A_{g_j}}\left(M_{g_j}, N_{g_j}/(IA_{g_j})^nN_{g_j}\right) \tag{14}$$

for all $i, n \geq 0$ and $j = 1, \ldots, m$. Therefore from (13) and (14), it is enough to prove the result for the ring $A_{g_j} = Q_j/\mathfrak{a}_j$ for each j. Note that Q_j is a regular ring of finite Krull dimension, and hence $\operatorname{projdim}_{Q_j}(M_{g_j})$ is finite. Therefore the result now follows by applying the Theorems 3.1 and 4.1 to each $A_{g_j} = Q_j/\mathfrak{a}_j$. □

Now we prove the analogous result (Theorem 5.2) in the projective case. Let K be a field not necessarily algebraically closed and $Q = K[X_0, X_1, \ldots, X_r]$ a polynomial ring over K, where $\deg(X_i) = 1$ for all i. Let $\mathfrak{a}$ be a homogeneous ideal of Q. Set $A = Q/\mathfrak{a}$. Suppose $A_{(\mathfrak{p})}$ is a complete intersection ring for every $\mathfrak{p} \in \operatorname{Proj}(A)$.

Let $\mathfrak{m}$ be the unique maximal homogeneous ideal of A. If E is a graded A-module, then define the set of relevant associated primes of E as

$$^*\operatorname{Ass}_A(E) = \operatorname{Ass}_A(E) \smallsetminus \{\mathfrak{m}\}.$$

With the above hypotheses we have the following theorem:

Theorem 5.2. *Let M, N be two finitely generated graded A-modules and I a homogeneous ideal of A. Then*

$$\bigcup_{i\geq 0}\bigcup_{n\geq 0} {}^*\operatorname{Ass}_A\left(\operatorname{Ext}^i_A(M, N/I^nN)\right) \quad \textit{is a finite set.}$$

Moreover, there exists $i_0, n_0 \geq 0$ such that for all $i \geq i_0, n \geq n_0$, we have

$$\begin{aligned} {}^*\operatorname{Ass}_A\left(\operatorname{Ext}^{2i}_A(M, N/I^nN)\right) &= {}^*\operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0}_A(M, N/I^{n_0}N)\right), \\ {}^*\operatorname{Ass}_A\left(\operatorname{Ext}^{2i+1}_A(M, N/I^nN)\right) &= {}^*\operatorname{Ass}_A\left(\operatorname{Ext}^{2i_0+1}_A(M, N/I^{n_0}N)\right). \end{aligned}$$

Proof. For a homogeneous element $x \in A$, set ${}^*D(x) = \{\mathfrak{p} \in \operatorname{Proj}(A) : x \notin \mathfrak{p}\}$. As in [7, *Proof of Theorem 6.3*], we have

$$\operatorname{Proj}(A) = {}^*D(g_1) \cup \cdots \cup {}^*D(g_m) \quad \text{for some homogeneous } g_1, \ldots, g_m \in A,$$

such that $A_{g_j} = Q_j/\mathfrak{a}_j$ for some regular ring Q_j of finite Krull dimension and some homogeneous ideal $\mathfrak{a}_j$ of Q_j generated by a Q_j-regular sequence. Therefore clearly for any graded A-module E, we have

$$^*\operatorname{Ass}_A(E) = \bigcup\{\mathfrak{q} \cap A : \mathfrak{q} \in \operatorname{Ass}_{A_{g_j}}(E_{g_j}) \text{ for some } j = 1, \ldots, m\}.$$

Similarly, as in the proof of Theorem 5.1, the result follows by applying the Theorems 3.1 and 4.1 to each $A_{g_j} = Q_j/\mathfrak{a}_j$. □

6 Support varieties

Let $(A, \mathfrak{m})$ be a local complete intersection ring of codimension c. Let M and N be two finitely generated A-modules. Define

$$\operatorname{cx}_A(M, N) = \inf\left\{b \in \mathbb{N} \;\middle|\; \limsup_{n\to\infty} \frac{\mu(\operatorname{Ext}^n_A(M, N))}{n^{b-1}} < \infty\right\},$$

where $\mu(D)$ denote the number of minimal generators of a finitely generated A-module D.

In this section, we prove the following theorem:

Theorem 6.1. *Let $(A, \mathfrak{m})$ be a local complete intersection ring. Let M, N be two finitely generated A-modules and I an ideal of A. Then*

$$\mathrm{cx}_A(M, N/I^jN) \quad \textit{is constant for all } j \gg 0.$$

As in [7, Section 7.2], we may assume that our local complete intersection A

(1) is complete. So $A = Q/(f_1, \ldots, f_c)$, where $(Q, \mathfrak{n})$ is a regular local ring and $\mathbf{f} = f_1, \ldots, f_c \in \mathfrak{n}^2$ is a Q-regular sequence.

(2) has an algebraically closed residue field k.

Let U, V be two finitely generated A-modules. Let

$$\mathrm{Ext}^\star_A(U, V) := \bigoplus_{i \geq 0} \mathrm{Ext}^i_A(U, V)$$

be the total Ext module of U, V over the ring $A[t_1, \ldots, t_c]$ of cohomology operators t_j defined by $\mathbf{f}$, where $\deg(t_j) = 2$ for all $j = 1, \ldots, c$. Set

$$\mathcal{C}(U, V) := \mathrm{Ext}^\star_A(U, V) \otimes_A k.$$

Since $\mathrm{Ext}^\star_A(U, V)$ is a finitely generated graded $A[t_1, \ldots, t_c]$-module, $\mathcal{C}(U, V)$ is a finitely generated graded module over

$$T := A[t_1, \ldots, t_c] \otimes_A k = k[t_1, \ldots, t_c].$$

Define the support variety $\mathscr{V}(U, V)$ of U, V as the zero set in k^c of the annihilator of $\mathcal{C}(U, V)$ in T, that is

$$\mathscr{V}(U, V) := \{(b_1, \ldots, b_c) \in k^c : P(b_1, \ldots, b_c) = 0, \ \forall\ P \in \mathrm{ann}_T(\mathcal{C}(U, V))\} \cup \{0\}.$$

By [1, 2.4.(2)], we have

$$\mathrm{cx}_A(U, V) = \dim(\mathscr{V}(U, V)) = \dim_T(\mathcal{C}(U, V)).$$

Before the proof of Theorem 6.1, we give the following well-known lemma:

Lemma 6.2. *Let $\mathscr{R} = \oplus_{n \geq 0} \mathscr{R}_n$ be a Noetherian standard $\mathbb{N}$-graded ring, $\mathscr{M} = \oplus_{n \geq 0} \mathscr{M}_n$ a finitely generated $\mathbb{N}$-graded $\mathscr{R}$-module. Then there exists $j_0 \geq 0$ such that*

$$\dim_{\mathscr{R}_0}(\mathscr{M}_j) = \dim_{\mathscr{R}_0}(\mathscr{M}_{j_0}) \quad \textit{for all } j \geq j_0.$$

Proof. Since $\mathscr{M}$ is a finitely generated $\mathbb{N}$-graded module over a Noetherian standard $\mathbb{N}$-graded ring $\mathscr{R}$, there exists $j_0 \geq 0$ such that

$$\mathrm{ann}_{\mathscr{R}_0}(\mathscr{M}_j) = \mathrm{ann}_{\mathscr{R}_0}(\mathscr{M}_{j_0}) \quad \text{for all } j \geq j_0,$$

and hence $\dim_{\mathscr{R}_0}(\mathscr{M}_j) = \dim_{\mathscr{R}_0}(\mathscr{M}_{j_0})$ for all $j \geq j_0$. □

6.3. Since $\mathrm{gr}_I(N) = \oplus_{j\geq 0} I^jN/I^{j+1}N$ is a finitely generated graded $\mathscr{R}(I)$-module, by Theorem 2.5,

$$\bigoplus_{j\geq 0} \mathrm{Ext}^\star_A(M, I^jN/I^{j+1}N) = \bigoplus_{i\geq 0}\bigoplus_{j\geq 0} \mathrm{Ext}^i_A(M, I^jN/I^{j+1}N)$$

is a finitely generated bigraded $\mathscr{R}(I)[t_1, \ldots, t_c]$-module, and hence

$$\bigoplus_{j\geq 0} \mathrm{Ext}^\star_A(M, I^jN/I^{j+1}N) \otimes_A k$$

is a finitely generated graded $\mathscr{R}(I)[t_1, \ldots, t_c] \otimes_A k = F(I)[t_1, \ldots, t_c]$-module, where $F(I)$ is the fiber cone of I which is a finitely generated k-algebra. Writing

$$F(I)[t_1, \ldots, t_c] = k[x_1, \ldots, x_m][t_1, \ldots, t_c] = T[x_1, \ldots, x_m],$$

and using Lemma 6.2, we have

$$\dim_T \left(\mathrm{Ext}^\star_A(M, I^jN/I^{j+1}N) \otimes_A k\right) \quad \text{is constant for all } j \gg 0,$$

and hence $\mathrm{cx}_A(M, I^jN/I^{j+1}N)$ is constant for all $j \gg 0$.

Now we give

Proof of Theorem 6.1. Fix $j \geq 0$. Consider the short exact sequence of A-modules

$$0 \longrightarrow I^jN/I^{j+1}N \longrightarrow N/I^{j+1}N \longrightarrow N/I^jN \longrightarrow 0,$$

which induces the following exact sequence of A-modules for each i:

$$\begin{aligned} &\mathrm{Ext}^{i-1}_A(M, N/I^jN) \to \\ &\mathrm{Ext}^i_A(M, I^jN/I^{j+1}N) \to \mathrm{Ext}^i_A(M,N/I^{j+1}N) \to \mathrm{Ext}^i_A(M, N/I^jN) \to \\ &\mathrm{Ext}^{i+1}_A(M, I^jN/I^{j+1}N). \end{aligned}$$

Taking direct sum over i and setting

$$U_j := \bigoplus_{i\geq 0} \mathrm{Ext}^i_A(M, I^jN/I^{j+1}N), \quad V_j := \bigoplus_{i\geq 0} \mathrm{Ext}^i_A(M, N/I^jN),$$

we have an exact sequence of $A[t_1, \ldots, t_c]$-modules:

$$V_j(-1) \xrightarrow{\varphi_1} U_j \xrightarrow{\varphi_2} V_{j+1} \xrightarrow{\varphi_3} V_j \xrightarrow{\varphi_4} U_j(1).$$

Set $Z_j := \mathrm{Image}(\varphi_1)$, $X_j := \mathrm{Image}(\varphi_2)$ and $Y_j := \mathrm{Image}(\varphi_3)$. Thus we have the following commutative diagram of exact sequences:

$$\begin{array}{ccccccccccc} V_j(-1) & \longrightarrow & U_j & \longrightarrow & V_{j+1} & \longrightarrow & V_j & \longrightarrow & U_j(1) \\ & \searrow & \uparrow & \searrow & \uparrow & \searrow & \uparrow & \searrow & \uparrow & & \\ & & Z_j & & X_j & & Y_j & & Z_j(1) & & \\ & \nearrow & & \searrow & \uparrow & \searrow & \uparrow & \searrow & \uparrow & \searrow & \\ 0 & & & & 0 & & 0 & & 0 & & 0. \end{array}$$

Now consider the short exact sequences of $A[t_1, \ldots, t_c]$-modules:

$$0 \to X_j \to V_{j+1} \to Y_j \to 0, \quad 0 \to Y_j \to V_j \to Z_j(1) \to 0.$$

Tensoring the above sequences with k over A, we get the following exact sequences of $T = k[t_1, \ldots, t_c]$-modules:

$$X_j \otimes_A k \xrightarrow{\Phi_j} V_{j+1} \otimes_A k \longrightarrow Y_j \otimes_A k \to 0,$$
$$Y_j \otimes_A k \xrightarrow{\Psi_j} V_j \otimes_A k \longrightarrow Z_j(1) \otimes_A k \to 0.$$

Now for each $j \geq 0$, set $X'_j := \text{Image}(\Phi_j)$ and $Y'_j := \text{Image}(\Psi_j)$ to get the following short exact sequences of T-modules:

$$0 \longrightarrow X'_j \longrightarrow V_{j+1} \otimes_A k \longrightarrow Y_j \otimes_A k \longrightarrow 0, \tag{15}$$
$$0 \longrightarrow Y'_j \longrightarrow V_j \otimes_A k \longrightarrow Z_j(1) \otimes_A k \longrightarrow 0. \tag{16}$$

From Section 6.3, $\oplus_{j\geq 0} U_j$ is a finitely generated graded $\mathscr{R}(I)[t_1, \ldots, t_c]$-module, and hence its submodule $\oplus_{j\geq 0} Z_j$ is also so. Therefore $\oplus_{j\geq 0}(Z_j \otimes_A k)$ is a finitely generated graded module over

$$\mathscr{R}(I)[t_1, \ldots, t_c] \otimes_A k = F(I)[t_1, \ldots, t_c] = T[x_1, \ldots, x_m].$$

Then by Lemma 6.2, $\dim_T(Z_j \otimes_A k) = z$ for all sufficiently large j, where z is some constant. Therefore considering the short exact sequences (15) and (16), we get that:

$$\dim_T(V_{j+1} \otimes_A k) = \max\{\dim_T(X'_j), \dim_T(Y_j \otimes_A k)\}, \tag{17}$$
$$\dim_T(V_j \otimes_A k) = \max\{\dim_T(Y'_j), z\} \geq z \tag{18}$$

for all sufficiently large j, say $j \geq j_0$.

Note that $\dim_T(V_j \otimes_A k) = \text{cx}_A(M, N/I^j N)$ for all $j \geq 0$. Thus it is enough to prove that the stability of $\dim_T(V_j \otimes_A k)$ holds for all sufficiently large j.

If $\dim_T(V_j \otimes_A k) = z$ for all $j \geq j_0$, then we are done. Otherwise there exists some $j \geq j_0$ such that $\dim_T(V_j \otimes_A k) > z$, and hence for this j we have

$$\dim_T(V_j \otimes_A k) = \dim_T(Y'_j) \leq \dim_T(Y_j \otimes_A k) \leq \dim_T(V_{j+1} \otimes_A k).$$

First equality above occurs from (18), second inequality occurs because Y'_j is a quotient module of $Y_j \otimes_A k$, and the last inequality occurs from (17).

Note that $\dim_T(V_{j+1} \otimes_A k) > z$. Then by applying the similar procedure, we have

$$\dim_T(V_{j+1} \otimes_A k) \leq \dim_T(V_{j+2} \otimes_A k).$$

In this way we have a bounded non-decreasing sequence

$$\dim_T(V_j \otimes_A k) \leq \dim_T(V_{j+1} \otimes_A k) \leq \dim_T(V_{j+2} \otimes_A k) \leq \cdots \leq \dim(T) < \infty,$$

which eventually stabilize somewhere, and hence the required stability holds. □

References

[1] L. L. Avramov and R.-O. Buchweitz. Support varieties and cohomology over complete intersections. *Invent. Math.* **142** (2000), 285-318.

[2] M. Brodmann. Asymptotic stability of $\mathrm{Ass}(M/I^nM)$. *Proc. Amer. Math. Soc.* **74** (1979), 16-18.

[3] D. Eisenbud. Homological algebra on a complete intersection, with an application to group representations. *Trans. Amer. Math. Soc.* **260** (1980), 35-64.

[4] T. H. Gulliksen. A change of ring theorem with applications to Poincaré series and intersection multiplicity. *Math. Scand.* **34** (1974), 167-183.

[5] D. Katz and E. West. A linear function associated to asymptotic prime divisors. *Proc. Amer. Math. Soc.* (6) **132** (2004), 1589-1597.

[6] L. Melkersson and P. Schenzel. Asymptotic prime ideals related to derived functors. *Proc. Amer. Math. Soc.* **117** (1993), 935-938.

[7] T. J. Puthenpurakal. On the finite generation of a family of Ext modules. *Pacific J. Math.* **266** (2013), 367-389.

[8] W. V. Vasconcelos. Cohomological degrees of graded modules. pp. 345-392 in Six lectures on commutative algebra (Bellaterra, 1996), edited by J. Elias et al., Progr. Math., **166**, Birkhäuser, Basel, 1998.

[9] E. West. Primes associated to multigraded modules. *J. Algebra* **271** (2004), 427-453.

Department of Mathematics
Indian Institute of Technology Bombay
Powai
Mumbai 400 076
India
e-mail: dipankar@math.iitb.ac.in
tputhen@math.iitb.ac.in